L Number	Hits	Search Text	DB	Time stamp
~~	1816	c ADJ elegans	USPAT;	2003/01/09 17:24
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
-	29	c ADJ elegans.clm.	USPAT;	2003/01/09 17:23
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
	321	c ADJ elegans and (phenoty\$5 SAME	USPAT;	2003/01/09 17:37
		screen\$5)	US-PGPUB;	
			EPO; JPO;	
	1.7	, and the second	DERWENT	0000/01/00 17 07
_	17	bogaert NEAR thierry	USPAT;	2003/01/09 17:37
			US-PGPUB;	
			EPO; JPO;	
	2	Foightinger MEAD wighows	DERWENT	2003/01/09 17:35
-		Feichtinger NEAR richard	USPAT; US-PGPUB;	2003/01/09 17:35
			EPO; JPO;	
			DERWENT	
_	1	kaletta NEAR titus	USPAT;	2003/01/09 17:40
	1	Naiceda MDMM Cleas	US-PGPUB;	2003/01/03 17.40
			EPO; JPO;	
,			DERWENT	
_	26	(constructing ADJ libraries) and elegans	USPAT;	2003/01/09 17:45
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
-	0	(constructing ADJ libraries) SAME elegans	USPAT;	2003/01/09 17:45
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
_	310	librar\$5 SAME elegans	USPAT;	2003/01/09 17:49
ĺ			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
-	168	leastone	USPAT;	2003/01/09 17:49
			US-PGPUB;	
			EPO; JPO;	
	1	1	DERWENT	0002/01/00 17 51
-	1	leastone and (nematode or elegans)	USPAT;	2003/01/09 17:51
			US-PGPUB;	
			EPO; JPO; DERWENT	
_	ے ا	leastone and gene	USPAT;	2003/01/09 17:51
	· ·	2000 cone and gone	US-PGPUB;	2000/01/00 17:01
			EPO; JPO;	
			DERWENT	
-	8	((US-6225120-\$ or US-6278039-\$ or	USPAT;	2003/01/09 17:59
		US-6329566-\$ or US-6433247-\$ or	US-PGPUB;	
		US-6465715-\$).did. or (US-20010016332-\$ or	EPO; JPO;	
		US-20020037585-\$ or US-20020064523-\$ or	DERWENT	
		US-20020194624-\$).did. or (WO-9638555-\$ or		
		WO-9824810-\$ or WO-9937770-\$ or		
		GB-2349217-\$ or WO-9964586-\$ or		
]	GB-2351496-\$ or GB-2358399-\$ or		
		GB-2358400-\$ or GB-2359358-\$).did. or		
		(GB-2359361-\$ or GB-2359358-\$ or		
		GB-2351152-\$).did.) and librar\$5		0000/07/77
-	10	Ruvkun NEAR Gary	USPAT;	2003/01/09 17:59
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	

_	24	(US-6225120-\$ or US-6278039-\$ or US-6329566-\$ or US-6433247-\$ or US-6465715-\$).did. or (US-20010016332-\$ or US-20020037585-\$ or US-20020064523-\$ or	USPAT; US-PGPUB; EPO; DERWENT	2003/01/09 18:02
		US-20020194624-\$).did. or (WO-9638555-\$ or WO-9824810-\$ or WO-9937770-\$ or GB-2349217-\$ or WO-9964586-\$ or GB-2351496-\$ or GB-2358399-\$ or		
		GB-2358400-\$ or GB-2359358-\$ or WO-9851351-\$ or WO-9630053-\$).did. or (GB-2359361-\$ or GB-2359358-\$ or GB-2351152-\$).did.		
_	13	US-6329566-\$ or US-6433247-\$ or US-6465715-\$).did. or (US-20010016332-\$ or US-20020037585-\$ or US-20020064523-\$ or	USPAT; US-PGPUB; EPO; DERWENT	2003/01/09 18:03
		US-20020194624-\$).did. or (WO-9638555-\$ or WO-9824810-\$ or WO-9937770-\$ or GB-2349217-\$ or WO-9964586-\$ or GB-2351496-\$ or GB-2358399-\$ or		
		GB-2358400-\$ or GB-2359358-\$ or WO-9851351-\$ or WO-9630053-\$).did. or (GB-2359361-\$ or GB-2359358-\$ or GB-2351152-\$).did.) and phenoty\$5		

Murakami et al, Genetics 143:1207-1218 1996.*

Strojek RM and Wagner TE Genetic Engineering: Principles and Methods 10:221-246, 1988.*

Larsan et al, Genetics 139:1567-1583, 1995.*

Kuo et al, PNAS, 92:6911-6914, 1995.*

Lonnqvist et al Nat. Med. 1(9):950-953, 1995.*

Galili et al., "Fusion of a fork head domain gene to PAX3 in the Solid Tumour Alveolar Rhabdomyosarcoma," Nat. Genet. 5:230-235 (1993).

Kimura et al., "daf-2, an Insulin Receptor-Like Gene That Regulates Longevity and Diapause in Caenorhabditis elegans," Science 277:942-946 (1997).

Lin et al., "daf-16: An HNF-3/Forkhead Family Member That Can Function to Double the Life-Span of Caenorhabditis elegans," Science 278:1319-1322 (1997).

McCombie et al., "Caenorhabditis elegans Expressed Sequence Tags Identify Gene Families and Potential Disease Gene Homologues," Nature Genetics 1:124-131 (1992).

Murakami et al., "A Genetic Pathway Conferring Life Extension and Resistance to UV Stress in Caenorhabditis elegans," Genetics 143:1207-1218 (1996).

Waterston et al., "A Survey of Expressed Genes in Caenorhabditis elegans," Nature Genetics 1:114-123 (1992).

Zwaal et al., "Target-Selected Gene Inactivation in Caenorhabditis elegans by Using a Frozen Transposon Insertion Mutant Bank," Proc. Natl. Acad. Sci. USA 90:7431-7435 (1993).

Gil et al., "Regulation of the Insulin-Like Developmental Pathway of Caenorhabditis elegans by a Homolog of the PTEN Tumor Suppressor Gene," Proc. Natl. Acad. Sci. USA 96:2925-2930 (1999).

Arpagaus, Vertebrate insulin induces diapause termination in Pieris brassicae pupae, Roux's Arch. Dev. Biol., 196:527-530 (1987).

Baker et al., A novel mesoderm inducer, Madr2, functions in the activin signal transduction pathway, Genes and Development, 10:1880-1889 (1996).

Bargmann et al., Control of Larval Development by Chemosensory Neurons in Caenorhabditis elegans, Science, 251:1243-1246 (1991).

Brown-Borg et al., Dwarf mice and the ageing process, Nature, 384:33 (1996).

Bruning et al., Development of a Novel Polygenic Model of NIDDM in Mice Heterozygous for IR and IRS-1 Null Alleles, Cell, 88:561-572 (1997).

Coleman, Obesity Genes: Beneficial Effects in Heterozygous Mice, Science, 203:663-665 (1979).

Dorman et al., The age-1 and daf-2 Genes Function in a Common Pathway to Control the Lifespan of Caenorhabditis elegans, Genetics, 141:1399-1406 (1995).

Ebina et al., The Human Insulin Receptor cDNA: The Structural Basis for Hormone-Activated Transmembrane Signalling, Cell, 40:747-758 (1985).

Estevez et al., The daf-4 gene encodes a bone morphogenetic protein receptor controlling C. elegans dauer larva development, Nature, 365:644-649 (1993).

Ewbank et al., Structural and Functional Conservation of the Caenorhabditis elegans Timing Gene clk-1, Science, 275:980-983 (1997).

Fernandez et al., The Drosophila insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential, EMBO J., 14:3373-3384 (1995).

Georgi et al., daf-1, a C. elegans Gene Controlling Dauer Larva Development, Encodes a Novel Receptor Protein Kinase, Cell, 61:635-645 (1990).

Golden et al., The Caenorhabditis elegans Dauer Larva: Developmental

Effects of Pheromone, Food and Temperature, Developmental Biology, 102:368-37 (1984).

Golden et al., A pheromone-induced developmental switch in Caenorhabditis elegans: Temperature-sensitive mutants reveal a wild-type temperature-dependent process, Proc. Natl. Proc. Acad. Sci. U.S.A., 81:819-823 (1984).

Gottlieb et al., daf-2, daf-16 and daf-23: Genetically Interacting Genes Controlling Dauer Formation in Caenorhabditis elegans, Genetics, 137:107-120 (1994).

Graff et al., Xenopus Mad Proteins Transduce Distinct Subsets of Signals for the TGF.beta. Superfamily, Cell, 85:479-487 (1996).

Green et al., Responses of Embryonic Xenopus Cells to Activin and FGF Are Separated by Multiple Dose Thresholds and Correspond to Distinct Axes of the Mesoderm, Cell, 71:731-739 (1992).

Hahn et al., DPC4, A Candidate Tumor Suppressor Gene at Human Chromosome 18q21.1, Science, 271:350-353 (1996).

Hemmings, Akt Signaling: Linking Membrane Events to Life and Death Decisions, Science, 275:628-630 (1997).

Hetru et al., Isolation and structural characterization of an insulin-related molecule, a predominant neuropeptide from Locusta migratoria, Eur. J. Biochem., 201:495-499 (1991).

Hoodless et al., MADR1, a MAD-Related Protein That Functions in BMP2 Signaling Pathways, Cell, 85:489-500 (1996).

Hotamisligil et al., Adipose Expression of Tumor Necrosis Factor-.alpha.: Direct Role in Obesity-Linked Insulin Resistance, Science, 259:87-91 (1993).

Hubbard et al., Crystal structure of the tyrosine kinase domain of the human insulin receptor, Nature, 372:746-754 (1994).

Jonas et al., Regulation by insulin of a unique neuronal Ca.sup.2+ pool and of neuropeptide secretion, Nature, 385:343-346 (1997).

Kahn et al., Genetics of Non-Insulin-Dependent (Type-II) Diabetes Mellitus, Annu. Rev. Med., 47:509-531 (1996).

Kawakami et al., Molecular Cloning of the Bombyx mori Prothoracicotropic Hormone, Science, 247:1333-1335 (1990).

Kenyon et al., A C. elegans mutant that lives twice as long as wild type, Nature, 366:461-464 (1993).

Kim et al., Detection of mutations in the insulin receptor gene in patients with insulin resistance by analysis of single-stranded conformational polymorphisms, Diabetologia, 35:261-266 (1992).

Kimble, Alterations in Cell Lineage following Laser Ablation of Cells in the Somatic Gonad of Caenorhabditis elegans, Developmental Biology, 87:286-300 (1981).

Klass, A Method for the Isolation of Longevity Mutants in the Nematode Caenorhabditis elegans and Initial Results, Mechanisms of Ageing and Dev., 22:279-286 (1983).

Krause, Transcription and Translation, Chapter 20, Methods Cell Biol., Academic Press, San Diego, CA, 48:483-512 (1995).

Lagna et al., Partnership between DPC4 and SMAD proteins in TGF-.beta. signalling pathways, Nature, 383:832-836 (1996).

Larsen et al., Genes that Regulate Both Development and Longevity in Caenorhabditis elegans, Genetics, 139:1567-1583 (1995).

Liu et al., A human Mad protein acting as a BMP-regulated transcriptional activator, Nature, 381:620-623 (1996).

Lonnqvist et al., Overexpression of the obese (ob) gene in adipose tissue of human obese subjects, Nat. Med. 1:950-953 (1995).

Macias-Silva et al., MADR2 Is a Substrate of the TGF.beta. Receptor and Its Phosphorylation Is Required for Nuclear Accumulation and Signaling, Cell, 87:1215-1224 (1996).

Malone et al., A Screen for Nonconditional Dauer-Constitutive Mutations in

Caenorhabditis elegans, Genetics, 136:879-886, (1994)

Mathews et al., Regulation of insulin-like growth factor I gene expression by growth hormone, Proc. Natl. Acad. Sci. U.S.A., 83:9343-9347 (1986).

Mello et al., Efficient gene transfer in C. elegans: extrachromosomal maintenance and integration of transforming sequences, EMBO J., 10:3959-3970 (1991).

Morris et al., A phosphatidylinositol-3-OH kinase family member regulating longevity and diapause in Caenorhabditis elegans, Nature, 382:536-539 (1996).

Nagasawa et al., Amino-Terminal Amino Acid Sequence of the Silkworm Prothoracicotropic Hormone: Homology with Insulin, Science, 266:1344-1345 (1984).

Ogg et al., The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in C. elegans, Nature, 389:994-999 (1997).

O'Riordan et al., Intermediary Metabolism in the Dauer Larva of the Nematode Caenorhabditis elegans-II. The Glyoxylate Cycle and Fatty-Acid Oxidation, Comp. Biochem. & Physiol., 95B:125-130 (1990).

O'Riordan et al., Intermediary Metabolism in the Dauer Larva of the Nematode Caenorhabditis elegans-l. Glycolysis, Gluconeogenesis, Oxidative Phosphorylation and the Tricarboxylic Acid Cycle, Comp. Biochem. & Physiol., 92B:233-238 (1989).

Popham et al., Aspects of the fine structure of the dauer larva of the nematode Caenorhabditis elegans, Can. J. Zool., 57:794-800 (1979).

Reinhardt et al., Selective Coexpression of Insulin Receptor-related Receptor (IRR) and TRK in NGF-Sensitive Neurons, J. Neurosci., 14:4674-4683 (1994).

Ren et al., Control of C. elegans Larval Development by Neuronal Expression of a TGF-beta. Homolog, Science, 274:1389-1391 (1996).

Riddle et al., Interacting genes in nematode dauer larva formation, Nature, 290:668-671 (1981).

Riddle, D. et al., Genetic and Environmental Regulation of Dauer Larva Develompent, C. elegans II, pp. 739-768 (1997).

Roovers et al., Characterization of a putative molluscan insulin-related peptide receptor, Gene, 162:181-188 (1995).

Savage et al., Caenorhabditis elegans genes sma-2, sma-3, and sma-4 define a conserved family of transforming growth factor .beta. pathway components, PNAS, 93:790-794 (1996).

Schackwitz et al., Chemosensory Neurons Function in Parallel to Mediate a Pheromone Response in C. clegans, Neuron, 17:719-728 (1996).

Shier et al., Primary Structure of a Putative Receptor for a Ligand of the Insulin Family, J. Biol. Chem., 264:14605-14608 (1989).

Songyang et al., SH2 Domains Recognize Specific Phosphopeptide Sequences, Cell, 72:767-778 (1993).

Swanson et al., Critical Periods in the Development of the Caenorhabditis elegans Dauer Larva, Developmental Biology, 84:27-40 (1981).

Taylor, Lilly Lecture: Molecular Mechanisms of Insulin Resistance, Diabetes, 41:1473-1490 (1992).

Thomas et al., Evidence for Parallel Processing of Sensory Information Controlling Dauer Formation in Caenorhabditis elegans, Genetics, 134:1105-1117 (1993).

Ullrich et al., Human insulin receptor and its relationship to the tyrosine kinase family of oncogenes, Nature, 313:756-761 (1985).

Ullrich et al., Insulin-like growth factor I receptor primary structure: comparison with insulin receptor suggests structural determinants that define functional specificity, EMBO J., 5:2503-2512 (1986).

Vowels et al., Genetic Analysis of Chemosensory Control of Dauer Formation in Caenorhabditis elegans, Genetics, 130:105-123 (1992).

Wadsworth et al., Developmental Regulation of Energy Metabolism in Caenorhabditis elegans, Developmental Regulation of Energy Metabolism in Developmental Regulation of Energy Metabolism in Caenorhabditis elegans, Developmental Regulation of Energy Metabolism in Caenorhabditis elegans in Caenorhabditis elegans elegans

White et al., The Insulin Signaling System, J. Biol. Chem., 269:1-4 (1994).

Wrana et al., MAD-related proteins in TGF-.beta. signalling, Trends Genet., 12:493-496 (1996).

Yoshimasa et al., Effects of Amino Acid Replacements within the Tetrabasic Cleavage Site on the Processing of the Human Insulin Receptor Precursor Expressed in Chinese Hamster Ovary Cells, J. Biol. Chem., 265:17230-17237 (1990).